



## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification <sup>6</sup> : C12N 15/06, C07K 16/22, C12N 5/20, G01N 33/577, 33/68, A61K 39/395, C07K 1/22 // 14/475	A1	(11) International Publication Number: <b>WO 96/30506</b> (43) International Publication Date: 3 October 1996 (03.10.96)
--	----	---

(21) International Application Number: PCT/JP96/00762

(22) International Filing Date: 22 March 1996 (22.03.96)

(30) Priority Data:  
7/65577 24 March 1995 (24.03.95) JP(71) Applicant (for all designated States except US): TAKEDA  
CHEMICAL INDUSTRIES, LTD. [JP/JP]; 1-1,  
Doshomachi 4-chome, Chuo-ku, Osaka-shi, Osaka  
541 (JP).


(72) Inventors; and

(75) Inventors/Applicants (for US only): SASADA, Reiko [JP/JP];  
6-8, Tenjin 4-chome, Nagaokakyo-shi, Kyoto 617 (JP).  
WATANABE, Tatsuya [JP/JP]; 50-1, Yamadaminami,  
Suita-shi, Osaka 565 (JP). TOYODA, Yukio [JP/JP];  
29-10-504, Minamimukonosono 8-chome, Amagasaki-shi,  
Hyogo 661 (JP).(74) Agents: OHTAWA, Akitoshi et al.; New Eiwa Building, 3-2,  
Nishishinbashi 2-chome, Minato-ku, Tokyo 105 (JP).(81) Designated States: AL, AM, AU, AZ, BB, BG, BR, BY, CA,  
CN, CZ, EE, GE, HU, IS, KE, KG, KR, KZ, LK, LR, LS,  
LT, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, RO,  
RU, SD, SG, SI, SK, TJ, TM, TR, TT, UA, UG, US, UZ,  
VN, European patent (AT, BE, CH, DE, DK, ES, FI, FR,  
GB, GR, IE, IT, LU, MC, NL, PT, SE).

Published

With international search report.

(54) Title: ANTIBODY TO HUMAN BETACELLULIN AND USE THEREOF



## (57) Abstract

Disclosed is an antibody which has a binding activity to human betacellulin protein or a mutein thereof with specificity; especially a monoclonal antibody which does not have cross reactivity with human epidermal growth factor (EGF) and human transforming growth factor  $\alpha$  (TGF- $\alpha$ ), belongs to the immunoglobulin class of IgG, and specifically binds to human betacellulin protein to neutralize biological activity thereof; a hybridoma for producing the monoclonal antibody; and a method for producing the monoclonal antibody. Said monoclonal antibody neutralizes biological activity of a human BTC protein, and binds to the protein with high sensitivity and specificity, so that they can be used as a therapeutic agent for diseases such as arterial sclerosis and cancers, and also used as a reagent for assaying the human BTC protein or a mutein thereof and as a diagnostic agent for diabetes or complications thereof.

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgyzstan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	KZ	Kazakhstan	SG	Singapore
CH	Switzerland	LI	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovakia
CM	Cameroon	LR	Liberia	SN	Senegal
CN	China	LT	Lithuania	SZ	Swaziland
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	LV	Latvia	TG	Togo
DE	Germany	MC	Monaco	TJ	Tajikistan
DK	Denmark	MD	Republic of Moldova	TT	Trinidad and Tobago
EE	Estonia	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	UG	Uganda
FI	Finland	MN	Mongolia	US	United States of America
FR	France	MR	Mauritania	UZ	Uzbekistan
GA	Gabon			VN	Viet Nam

## ANTIBODY TO HUMAN BETACELLULIN AND USE THEREOF

FIELD OF THE INVENTION

The present invention relates to an antibody to a human betacellulin (hereinafter also referred to as "BTC") protein or a mutein thereof, a hybridoma, their production and use thereof.

BACKGROUND OF THE INVENTION

As to the family of epidermal growth factors, since epidermal growth factor (EGF) was discovered by S. Cohen, 10 J. Biol. Chem., 237, 1555 (1962), Dev. Biol., 12, 394 (1965), various factors have been discovered. These factors are considered to have various functions such as differentiation, maturation, survival, functional retention and proliferation of not only epidermal cells, but also of 15 various cells. Specifically, these factors include transforming growth factor  $\alpha$  (TGF- $\alpha$ ) [G. J. Todaro and J. E. DeLarco, Nature, 264, 26 (1976) and H. Marquardt et al., Proc. Natl. Acad. Sci. USA, 80, 4684 (1983)], amphiregulin [M. Shoyab et al., Science, 243, 1074 (1989)] and heparin- 20 binding EGF (HB-EGF) [S. Higashiyama et al., Science, 251, 936 (1991)], in addition to EGF described above.

Human BTC protein is described in EP-A 0555785, and reported in Biochem. Biophys. Res. Commun., 190, 1173-1179 (1993). BTC protein was discovered as a novel cell growth 25 factor produced by a transgenic mouse-derived pancreatic beta tumor cell [Y. Shing et al., Science, 259, 1604 (1993)]. Further, based on the nucleotide sequence of the

- 2 -

gene thereof, the human BTC gene was cloned, and the characteristics of BTC protein were studied. As a result, (1) a precursor molecule having a transmembrane domain is processed at the C-terminus to produce matured type BTC protein, (2) the BTC gene is expressed in the normal mouse kidney, lung, liver and pancreatic beta cells, (3) BTC protein exhibits growth promoting activity on a vascular smooth muscle cell and a retinal pigment epithelial cell, and (4) BTC protein acts on an EGF receptor as a ligand [T. Watanabe et al., J. Biol. Chem., 269, 9966 (1994)] were revealed. From these, it is conceivable that BTC protein plays an important role in various organs, and abnormal expression thereof possibly induces diseases. Further, these results suggest the possibility that BTC protein contributes to the growth of smooth muscle cells relating to the crisis and progress of arterial sclerosis, and particularly to the possibility that BTC protein is involved in the crisis of diabetic vascular complications such as diabetic arterial sclerosis and diabetic retinitis or in the canceration of cells.

An antibody inhibiting BTC protein activity is considered to be used as one therapeutic agent for diseases in which BTC protein is involved. However, such a report which relates to an antibody to a human BTC protein has not been presented till now. The assay of the human BTC protein level in the body fluids is also considered as one means for diagnosing the diseases as described above.

- 3 -

However, it has not yet been achieved.

As described above, human BTC protein may participate in the crisis and progress of arterial sclerosis, and particularly in the crisis of diabetic vascular complications or the canceration of cells by its abnormal expression (excess expression). Accordingly, inhibition of excess activity of human BTC protein can conceivably treat these diseases. Therefore, the measurement of human BTC protein level in blood makes it possible to diagnose these diseases. In order to assay human BTC protein existing only in slight amounts in the body fluids, an assay with high sensitivity is required.

#### SUMMARY OF THE INVENTION

In view of the situations described above, the present inventors have prepared antibodies which offer an assay with high sensitivity for a human BTC protein and which are capable of neutralizing human BTC protein activity. As a result of further intensive investigation based thereon, the present inventors have completed the present invention.

In accordance with the present invention, there are provided

- (1) An antibody which specifically binds to human BTC protein or a mutein thereof;
- (2) The antibody according to (1) wherein human BTC protein has an amino acid sequence of SEQ ID NO:1;
- (3) The antibody according to (1) or (2) which is a monoclonal antibody;

- 4 -

(4) The antibody according to anyone of (1) to (3) which has no cross-reactivity with at least one of human epidermal growth factor, human transforming growth factor  $\alpha$  and mouse BTC protein;

5 (5) A monoclonal antibody which specifically binds to human BTC protein and is capable of neutralizing the biological activity of said protein, which does not have cross-reactivity with human epidermal growth factor, human transforming growth factor  $\alpha$  and mouse BTC protein, and  
10 which belongs to the immuno-globulin class of IgG;

(6) The antibody according to anyone of (1) to (5) which recognizes an amino acid sequence of 31st (Arg) to 80th (Tyr) of SEQ ID NO:1;

(7) A cloned hybridoma which is composed of a mammalian  
15 spleen cell immunized by human BTC protein or a mutein thereof and a homogenic or heterogenic lymphocyte;

(8) A method of producing the monoclonal antibody of (5), which comprises proliferating the hybridoma according to (7) in a liquid culture medium or in an abdominal cavity of  
20 a mammal to form and accumulate the monoclonal antibody and then collecting the monoclonal antibody;

(9) A method of detecting and assaying a human BTC protein or a mutein thereof which comprises contacting the antibody according to anyone of (1) to (6) with a specimen;

25 (10) A method of diagnosing diabetes or complications thereof which comprises contacting the antibody according to anyone of (1) to (6) with a specimen and assaying a

- 5 -

human BTC protein or a mutein thereof in the specimen;

(11) A pharmaceutical composition which comprises an effective amount of the antibody according to anyone of (1) to (6) and a pharmaceutically acceptable carrier, excipient or diluent;

(12) The pharmaceutical composition according to (11) for diagnosing diabetes or complications thereof;

(13) A kit for diagnosing diabetes or complications thereof which comprises an effective amount of the antibody according to anyone of (1) to (6);

(14) A kit for detecting and assaying a human BTC protein or a mutein thereof which which comprises an effective amount of the antibody according to anyone of (1) to (6); and

(15) A method of purifying a human BTC protein or a mutein thereof which comprises contacting the antibody according to anyone of (1) to (6) with a crude sample containing the protein or a mutein thereof and isolating it.

#### BRIEF DESCRIPTION OF THE DRAWINGS

Fig. 1 is a schematic representation showing the construction of plasmid pTB1515;

Fig. 2 is a schematic representation showing the construction of plasmid pTB1507;

Figs. 3-1 and 3-2 show a nucleotide sequence of cDNA of BTC-I obtained in Example 4 of EP-A 0555785 and an amino acid sequence deduced therefrom;

Fig. 4 is a schematic representation showing the

construction of plasmid pTB1516;

Fig. 5 is a graph showing results of S-Sepharose column chromatography and DNA synthesis induction activity obtained in Reference Example 5;

5 Fig. 6 is a graph showing results of gel filtration obtained in Reference Example 5;

Fig. 7 is a graph showing results of heparin high performance liquid column chromatography obtained in Reference Example 5;

10 Fig. 8 is a graph showing results of reverse high performance liquid chromatography obtained in Reference Example 5;

Fig. 9 shows results of SDS-PAGE silver staining obtained in Reference Example 5;

15 Fig. 10 is a graph showing results of S-Sepharose column chromatography obtained in Reference Example 6;

Fig. 11 is a graph showing results of reverse high performance liquid column chromatography obtained in Reference Example 6;

20 Fig. 12 shows results of SDS-PAGE silver staining obtained in Reference Example 6;

Fig. 13 is a graph showing results of reverse high performance liquid column chromatography obtained in Reference Example 6;

25 Fig. 14 shows N-terminal amino acid sequences of BTC-I and BTC-II obtained in Reference Example 6;

Fig. 15 is a graph showing the neutralizing activity



- 7 -

of a monoclonal antibody of the invention on human BTC in Example 7;

Fig. 16 is a graph showing a detection curve of human BTC;

5 Fig. 17 shows results of Coomassie Blue staining obtained in Example 9;

Fig. 18 is a graph showing results of reverse high performance liquid column chromatography obtained in Example 9; and

10 Fig. 19 shows results of silver staining obtained in Example 9.

#### DESCRIPTION OF THE PREFERRED EMBODIMENTS

Human BTC proteins are targets of the antibodies of  
15 the present invention and may be any as long as they are human-derived proteins having BTC-like activity, namely cell growth stimulating activity, such as fibroblasts, vascular smooth muscle cells and retinal pigment epithelial cells. They may be natural proteins extracted from various  
20 organs such as a liver, a kidney, etc. or recombinant proteins produced by genetic engineering technique.

Examples of the human-derived BTC proteins which can be used include a mature type protein having the amino acid sequence represented by SEQ ID NO:1 (refer to EP-A 0555785)  
25 and a precursor protein having the amino acid sequence represented by SEQ ID NO: 2.

These BTC proteins may be simple proteins consisting

- 8 -

of amino acids alone or conjugated proteins such as glycoproteins, lipoproteins, hemoproteins, metalloproteins, flavoproteins and phosphoproteins. In the case of glycoproteins, sugar chains include neutral sugars such as  
5 D-mannose, D-galactose and L-fructose, amino sugars such as D-glucosamine and D-galactosamine, and sialic acid.

Muteins of the human BTC proteins which can be used include, for example, deletion type muteins in which at least one constituent amino acid is deleted from each of  
10 the above-mentioned human BTC proteins, substitution type muteins in which at least one constituent amino acid of each of the BTC proteins is substituted by another amino acid(s), and addition type muteins in which at least one constituent amino acid is added to each of the BTC  
15 proteins. The numbers of deleted, substituted or added amino acids may be any as long as the inherent activities of the BTC proteins are not lost. Specifically, there can be used proteins having (1) amino acid sequences in which about 1 to about 40 amino acids are deleted from the amino  
20 acid sequence represented by SEQ ID NO: 1 or SEQ ID NO: 2, (2) amino acid sequences in which about 1 to about 10 amino acids of the amino acid sequence represented by SEQ ID NO: 1 or SEQ ID NO: 2 are substituted by another amino acid(s) or (3) amino acid sequences in which about 1 to about 40  
25 amino acids are added to the amino acid sequence represented by SEQ ID NO: 1 or SEQ ID NO: 2.

Of these muteins, deletion type muteins are preferred.

- 9 -

For example, proteins having amino acid sequences in which about 1 to about 40 amino acids are deleted from the N-terminus of the amino acid sequence represented by SEQ ID NO: 1 are preferably used. More specifically, a deletion  
5 type mutein of the BTC protein having an amino acid sequence in which 12 or 30 amino acids are deleted from the N-terminus of the amino acid sequence represented by SEQ ID NO: 1 are used [Watanabe et al., Journal of Biochemistry, 269, 9966 (1994)].

10 Other muteins of the BTC proteins may be used as long as the inherent biological activities of the BTC proteins are not lost. Examples of such muteins include muteins in which the N-termini of the BTC proteins are acylated, glycosylated, or chemically modified with polyethylene  
15 glycol derivatives.

In particular, in the present invention, human BTC protein having the amino acid sequence of SEQ ID NO:1 (BTC-I) is preferably used as the human BTC protein. In this case, the protein may be a mixture of one further having a  
20 methionine residue (Met) at the amino terminus thereof and one having no methionine residue, or may be one having no methionine residue (Met) at the amino terminus and starting with the subsequent aspartic acid residue (Asp).

As the mutein of the BTC protein, there is preferably  
25 used a deletion type human BTC mutein (human BTC-II) in which 30 amino acids are deleted from the N-terminus of the amino acid sequence of BTC-I having the amino acid sequence

- 10 -

represented by SEQ ID NO: 1.

Of the BTC proteins and the muteins thereof used in the present invention, the human BTC protein having the amino acid sequence represented by SEQ ID NO: 1 and the  
5 deletion type human BTC mutein having the amino acid sequence in which 12 or 30 amino acids are deleted from the N-terminus of the amino acid sequence represented by SEQ ID NO: 1 are known. The other BTC proteins and the muteins thereof can be prepared by methods known in the art or  
10 methods based thereon.

Methods of preparing the antibodies of the present invention are shown as follows.

The monoclonal antibodies of the present invention which specifically bind to a human BTC protein or a mutein  
15 thereof and neutralize biological activity thereof are prepared by immunizing mammals with a BTC protein or a mutein thereof. When the mammals are immunized with a human BTC protein or a mutein thereof described above, the mammals used for immunization include experimental animals  
20 such as sheep, goat, rabbits, guinea pigs, rats and mice. In order to obtain the monoclonal antibodies, rats and mice are preferred, and mice are particularly suitable. When mice are immunized, for example, any of the subcutaneous, intraperitoneal, intravenous, intramuscular and  
25 intracutaneous routes is available. Mainly, subcutaneous, intraperitoneal and intravenous injections are preferred, and the subcutaneous injection is particularly preferred.

- 11 -

The immunizing interval and the immunizing dose are widely changeable, and various methods are available. For example, immunization is generally carried out about 2 to about 6 times at intervals of about 2 weeks and spleen cells removed about 1 to about 5 days, preferably about 2 to about 4 days after the final immunization. The immunizing dose is preferably about 0.1  $\mu$ g or more, and more preferably about 10 to about 300  $\mu$ g per mouse, as the peptide amount per one immunization. Further, it is preferred to carry out a cell fusion using the spleen cells after confirmation of an increase in antibody titer in the blood by collecting partial blood before removal of the spleens.

The thus-prepared spleen cells are fused with lymphoid cells. For example, the spleen cells removed from the mice are fused with lymphoid cells such as homogeneous or heterogeneous (preferably homogeneous) myeloma cells, for example, P3-X63-Ag-8UI [Ichimori et al., J. Immun. Method, 80, 55 (1985)], having characteristics useful as selection marker such as hypoxanthine-guanine-phosphoribosyl transferase deficient (HGPRT<sup>-</sup>) or thymidine kinase deficient (TK<sup>-</sup>). The hybridoma cells can be produced by fusion, for example, in accordance with the method of Köhler and Milstein [Nature, 256, 495 (1975)]. Namely, for example, the myeloma cells and the spleen cells are suspended in a ratio of about 1:5 in a 1:1 mixed medium of Iscove medium and Ham F-12 medium (hereinafter referred to

- 12 -

as IH medium), and a fusogen such as Sendai virus or polyethylene glycol (PEG) is added thereto. It is of course possible to add another fusogen such as dimethyl sulfoxide (DMSO). The molecular weight of PEG is usually  
5 about 1,000 to about 6,000, time for the fusion is from about 0.5 to about 30 minutes, and the concentration is from about 10 to about 80%. As a preferred example, fusion is conducted in a concentration of from about 35 to about 55% for from about 4 to about 10 minutes using PEG 6,000,  
10 which results in efficient fusion. The fused cells can be selectively proliferated using hypoxanthine-aminopterin-thymidine medium (HAT medium) [Nature, 256, 495 (1975)].

The culture supernatant of the proliferated cells can be screened whether a desired antibody has been produced or  
15 not. The screening of the antibody titer can be carried out in the following manner. In this case, first, the presence or absence of the antibody produced by peptide immunization can be examined by known assays such as radio-immunoassays (RIAs) or enzyme immunoassays (EIAs). For  
20 these methods, various modified assays are also available. As a preferred example of the assays, a method using the EIA is hereinafter described. A rabbit anti-mouse immunoglobulin antibody is coupled to a carrier such as cellulose beads according to conventional methods, and then  
25 a culture supernatant or mouse serum to be assayed is added thereto, followed by reaction at a room temperature (about 4 to about 40°C in this specification, hereinafter the

- 13 -

same) for a defined time. After the reaction product is thoroughly washed, an enzyme-labeled peptide obtained by chemically conjugating the enzyme with the peptide and purifying the resulting product is added thereto and the  
5 reaction is carried out at a room temperature for a defined time. After the reaction product is thoroughly washed, an enzyme substrate is added thereto, followed by reaction at a room temperature for a defined time. Then, the absorbance or fluorescence of the product can be measured.

10 The cells in wells which show cell proliferation in a selective medium and antibody activity upon the peptide used for immunization is preferably cloned by a limiting dilution method. The supernatant of the cloned cells is screened for a high antibody titer, proliferating cells in  
15 a well which shows higher antibody titer, thereby obtaining monoclonal antibody-producing hybridoma clones showing reactivity with the immunized peptide.

The hybridoma cells thus cloned are proliferated, for example, in a liquid medium. Specifically, for example,  
20 the hybridoma cells are cultivated in a known liquid medium such as a medium comprising RPMI-1640 [G. E. Moore et al., J. Am. Med. Assoc., 199, 549 (1967)] with about 0.1-40% bovine serum for about 2 to about 10 days, preferably for about 3 to about 5 days, whereby the monoclonal antibody  
25 can be obtained from the cultivate solution. The desired antibody can also be obtained by intraperitoneally inoculating mammals with the above-mentioned monoclonal

- 14 -

antibody-producing hybridoma cells to proliferate the hybridoma cells, and then collecting the ascites. For example, in the case of mice, about  $1 \times 10^4$  to about  $1 \times 10^7$  cells, preferably about  $5 \times 10^5$  to about  $2 \times 10^6$  cells of the above-mentioned monoclonal antibody-producing hybridoma are intraperitoneally inoculated into mice such as BALB/c previously inoculated with mineral oil, etc., and the ascites is collected after about 7 to about 20 days, preferably after about 10 to about 14 days. From the antibodies produced and accumulated in the ascites, the desired BTC monoclonal antibody can be easily isolated as pure immunoglobulin by ammonium sulfate fractionation, DEAE-cellulose column chromatography, etc.

The polyclonal antibodies of the present invention can be prepared by methods known in the art or methods based thereon.

Immunogens used herein include, for example, the human BTC polypeptides, the muteins thereof and polypeptides having parts of the amino acid sequence from the 31st (Arg) to the 80th (Thr) of the amino acid sequence represented by SEQ ID NO: 1. The polypeptides used as an antigen are preferably the polypeptides having more than 10 amino acid residues. For example, the polypeptide having 15 amino acid residues from 66th to 88th of the amino acid sequence represented by SEQ ID NO:1 can be used as an antigen. The polypeptides can be prepared by peptide synthesis methods known in the art or methods based thereon, and either the



- 15 -

solid phase synthesis methods or the liquid phase synthesis methods may be employed.

The polyclonal antibodies can be prepared by producing conjugates of the immunogens and carrier polypeptides, immunizing mammals with the conjugates in a manner similar to that of the above-mentioned monoclonal antibodies, collecting products containing antibodies of BTC or muteins thereof from the immunized mammals, and isolating and purifying the antibodies.

10 As to the conjugates of the immunogens and the carrier proteins used for immunizing the mammals, the kinds of carrier proteins and the mixing ratios of the carrier proteins and the haptens may be any, as long as the antibodies can be efficiently obtained to the haptens coupled to the carrier proteins for immunization. For example, bovine serum albumin, bovine thyroglobulin or hemocyanin is coupled to the hapten in a weight ratio of about 0.1 to about 20:1, preferably about 1 to about 5:1.

20 In coupling of the haptens and the carrier proteins, various coupling agents can be used. Examples of the coupling agents include glutaraldehyde, carbodiimide reagents, maleimide active ester reagents and active ester reagents having thiol or dithiopyridyl groups.

25 The condensate is given to a mammal at a site where an antibody may be produced by administration of the condensate, alone or together with a carrier and/or a diluent. In order to enhance antibody production ability,

- 16 -

Freund's complete adjuvant or Freund's incomplete adjuvant may be given. The condensate is usually given once for every 2 to 6 weeks, totally about 3 to about 10 times. The mammals used for immunization include monkeys, rabbits, dogs, guinea pigs, mice, rats, sheep and goat.

The antibodies are collected, for example, from the blood or the ascites of the immunized mammals, and preferably from the blood thereof.

The anti-BTC antibody titer in the antisera can be assayed in the same manner as with the above-mentioned assay of the antibody titer in the hybridoma cultivate supernatants. Isolation and purification of the antibodies are performed according to the isolation and purification methods of immunoglobulins similarly to those of the above-mentioned monoclonal antibodies.

The antibodies of the present invention bind to various types of human BTC proteins or muteins thereof described above with high sensitivity and specificity.

The antibodies of the present invention which further have the following characteristics in addition to the characteristics described above are preferred.

a) They do not neutralize biological activity of mouse BTC protein, but neutralize biological activity of human BTC.

b) They do not have immune cross-reactivity with human TGF- $\alpha$  and human EGF.

c) They belong to the immunoglobulin class of IgG<sub>1</sub>

- 17 -

subclass, and have a molecular weight of 140,000 to 160,000.

d) They recognize specifically the amino acid sequence from 31st (Arg) to 80th (Tyr) of SEQ ID NO:1 (50-amino acid sequence).

Examples of the mouse BTC proteins used include a maturation type mouse BTC protein having the amino acid sequence represented by SEQ ID NO: 3 [Y. Shing et al., Science, 259, 1604 (1993)]. The amino acid sequence of this mature type mouse BTC protein has about 80% homology with that of the mature type human BTC protein having the amino acid sequence represented by SEQ ID NO: 1.

However, the preferred antibodies of the present invention, especially the monoclonal antibodies of the present invention, have the very unique characteristics that they do not bind to the mouse BTC proteins, but specifically bind to the human BTC proteins.

For example, the human BTC monoclonal antibodies produced from the hybridoma obtained in the example described below have the above-mentioned characteristics (a) to (d).

The antibodies of the present invention may be any, as long as they specifically bind to human BTC and neutralize biological activity thereof. The biological activities of the human BTC protein neutralized by the antibody of the present invention are, for example, cell growth stimulating activity such as fibroblasts, vascular smooth muscle cells

or retinal pigment epithelial cells. Examples of the monoclonal antibodies include antibodies 1A1, 2B2 and 5E5 produced by anti-human BTC monoclonal antibody-producing hybridomas 1A1, 2B2 and 5E5 obtained in the example 2 described below.

Examples of the polyclonal antibodies include one obtained in the example 10 described below.

For example, the monoclonal antibodies obtained in the examples described below inhibit a DNA synthesis induction of resting mouse 3T3 cells in the presence of 1 ng/ml of human BTC by addition of 1 µg/ml of the antibodies. Further, 10 pg/ml of a human BTC protein can be detected by a sandwich ELISA method using the monoclonal antibody and a biotinated antibody.

The antibodies of the present invention binding to a human BTC protein or a mutein thereof can be also used for detection and determination of a human BTC protein or a mutein thereof.

Detection and determination of a human BTC protein or a mutein thereof using the antibodies of the present invention can be performed, for example, by the immunochemical assay of measuring the human BTC protein using an human BTC antibody of the present invention bound to a carrier and a conjugate in which a labeling agent is directly bound to an anti-BTC antibody different from the former antibody in the antigen recognition site. The enzyme immunoassay is preferably employed among others.

The antibodies of the present invention are used as both the antibody held on the carrier and the antibody different from the above-mentioned antibody in the antigen recognition site used in the above-mentioned assay, and the 5 monoclonal antibodies are preferably used. Preferred examples of the monoclonal antibodies include antibodies 1A1, 2B2 and 5E5 prepared in the example described below, and they may be appropriately selected.

Examples of the carriers to which the antibodies are 10 bound in the above-mentioned assays include gel particles for example, agarose gels such as Sepharose 4B and Sepharose 6B (Pharmacia Fine Chemical, Sweden), dextran gels such as Sephadex G-75, Sephadex G-100 and Sephadex G-200 (Pharmacia Fine Chemical, Sweden) and polyacrylamide 15 gels such as Biogel P-30, Biogel P-60 and Biogel P-100 (Bio RAD Laboratories, U.S.A.); cellulose particles, for example, Avicel (Asahi Chemical Industry, Japan) and ion exchange cellulose such as diethylaminoethyl cellulose and carboxymethyl cellulose; physical adsorbents, for example, 20 glass such as glass balls, glass rods and aminoalkyl glass rods, silicone pieces, styrenic resins such as polystyrene balls and polystyrene particles, and plates for immunoassay (for example, Nunc, Denmark); and ion exchange resins, for example, weakly acidic cation exchange resins such as 25 Amberlite IRC-50 (Rohm & Haas, U.S.A.) and Zeocurve 226 (Permutit, West Germany), and weakly alkalic anion exchange resins such as Amberlite IR-4B and Dowex (Dow Chemical,

U.S.A.).

In order to bind the antibody on the carrier, conventional methods can be used. Examples of such methods include the cyanogen bromide method and the glutaraldehyde method which are described in Metabolism, 8, 696 (1971). As a simpler method, the antibody may be physically adsorbed on the surface of the carrier.

Examples of the labeling agents in the labeling agent-bound antibody conjugates include radioisotopes, enzymes, fluorescent substances and luminous substances, and enzymes are preferably used. Enzymes which are stable and high in specific activity are preferred, and peroxidases, alkaline phosphatases,  $\beta$ -D-galactosidases, glucose oxidases and the like can be used. Peroxidases are preferred among others, and peroxidases of various origins can be used. Examples thereof include peroxidases derived from horseradishes, pineapples, figs, sweet potatoes, broad beans and corn. Horseradish peroxidase (HRP) extracted from horseradishes is preferred among others.

In binding the peroxidase to the antibody, the peroxidase into which a maleimido group is preliminarily introduced is conveniently used to utilize the thiol group of Fab' as an antibody molecule.

When the maleimido group is introduced into the peroxidase, the maleimido group can be introduced through an amino group of the peroxidase. For this purpose, N-succinimidyl-maleimide-carboxylate derivatives can be used,

and N-( $\gamma$ -maleimidobutyloxy)succinimide (hereinafter also referred to as "GMBS") is preferably used. A certain group may therefore intervene between the maleimide group and the peroxidase.

5        GMBS reacts with the peroxidase in a buffer having a pH of about 6 to about 8 at about 10 to about 50°C for about 10 minutes to about 24 hours. The buffer solutions include, for example, 0.1 M phosphate buffer (pH 7.0). The maleimidated peroxidase thus prepared can be purified, for  
10    example, by gel chromatography. Examples of carriers used in the gel chromatography include Sephadex G-25 (Pharmacia Fine Chemical, Sweden) and Biogel P-2 (Bio RAD Laboratories, U.S.A.).

      The maleimidated peroxidase can be reacted with the  
15    antibody molecule in a buffer at about 0 to about 40°C for about 1 to about 40 hours. Examples of the buffers include 0.1 M phosphate buffer (pH 6.0) containing 5 mM sodium ethylenediaminetetraacetate. The peroxidase-labeled antibody thus prepared can be purified, for example, by gel  
20    chromatography. Examples of carriers used in the gel chromatography include Sephadex G-25 (Pharmacia Fine Chemical, Sweden) and Biogel P-2 (Bio RAD Laboratories, U.S.A.).

      Further, a thiol group may be introduced into the  
25    peroxidase to react with the maleimidated antibody molecule.

      Enzymes other than the peroxidases can also be

- 22 -

directly bound to antibodies based on the methods used for binding the peroxidases. Known methods which achieve such binding include, for example, the glutaraldehyde method, the periodic acid method and the water-soluble carbodiimide  
5 method.

Specimens to be tested in an assay system (detection and determination of human BTC protein) used in the present invention include the body fluids such as the urine, the serum, the plasma and the cerebrospinal fluid, extract of  
10 animal cells, and culture supernatants thereof.

As an example of assays (detection and determination of human BTC protein) used in the present invention, a case is hereinafter described in detail in which a peroxidase is used as the labeling agent, but the present invention is  
15 not limited thereto.

(1) First, a specimen containing a human BTC protein to be assayed is added to an antibody bound to a carrier to conduct the antigen-antibody reaction, and then the conjugate of the peroxidase with the anti-human BTC protein  
20 antibody obtained above is added thereto to allow them to react with each other.

(2) A substrate of the peroxidase is added to the reaction product obtained in (1), and then the absorbance or the fluorescent intensity of the resulting substance is  
25 measured, thereby detecting enzyme activity of the above-mentioned reaction product.

(3) A standard solution containing a known amount of



- 23 -

the human BTC protein is previously subjected to the procedures of the above (1) and (2) to prepare a standard curve showing the relation between the amount of the human BTC protein and the absorbance or the fluorescent intensity thereof.

(4) The absorbance or the fluorescent intensity obtained for the specimen (sample to be tested) containing an unknown amount of the human BTC protein is applied to the standard curve to determine the amount of the human BTC protein contained in the specimen.

This method of detecting and assaying a human BTC protein or a mutein thereof can be used for a diagnosis of diabetes or complications thereof.

The antibodies of the present invention can be used for a purification of a human BTC protein or a mutein thereof in a crude sample.

For the purification of the human BTC protein, the purified antibody of the present invention is coupled with an appropriate carrier such as activated agarose gel beads according to conventional methods, followed by packing in a column. Then, a sample containing the crude human BTC protein, such as a culture supernatant or a fluid of disrupted cells, is loaded onto the antibody column to allow the sample to be adsorbed thereby, followed by washing. Thereafter, the column is eluted with a chaotropic reagent such as potassium thiocyanate (KSCN) or under such acidic conditions that human BTC is not

- 24 -

inactivated. Thus, the human BTC protein can be efficiently purified.

The antibody column using the antibodies of the present invention can be prepared by coupling the  
5 monoclonal antibody of the present invention with an appropriate carrier, said antibody being, for example, purified from ascites or other fluids inoculated with hybridoma cells, in the following manner.

Any carrier may be used as long as the human BTC  
10 protein is specifically efficiently adsorbed thereby after coupling and suitable elution is thereafter possible. Examples of the carriers include agarose, cellulose and acrylamide polymers. As an example, agarose gel beads in which a primary amine of a protein is activated so as to be  
15 easily bindable, such as Affi-Gel 10 (Bio RAD), are conveniently used according to the following method. The antibody is reacted with Affi-Gel 10 in a buffer such as a bicarbonate solution having a concentration of about 0.001 to about 1 M, preferably about 0.1 M. The reaction can be  
20 conducted at about 0 to about 20°C at various pH values for about 10 minutes to about 24 hours, and preferably at about 4°C at a pH of about 3 to about 10 for about 4 hours. With respect to the mixing ratio of the antibody to Affi-Gel 10, the amount of the antibody which becomes bound to Affi-Gel  
25 10 increases as the amount of the antibody mixed therewith increases, until the ratio reaches about 50 mg of antibody per 1 ml of Affi-Gel 10. Hence, any ratio can be employed

- 25 -

within this range. However, about 10 to about 30 mg of the antibody is conveniently used, considering the binding efficiency and the purification efficiency in affinity column chromatography. The antibody-carrier conjugate thus  
5 formed is thoroughly washed with the buffer used in the reaction. Then, residual unreacted active groups are blocked by allowing the washed conjugate to stand for several days, or by adding a compound containing a primary amine such as ethanolamine-hydrochloric acid or glycine  
10 thereto to a final concentration of about 0.05 to about 0.10 M, followed by reaction at about 4°C for about 1 to about 4 hours, or by reacting a protein such as 1-5% bovine serum albumin (BSA) therewith at 4°C overnight. The conjugate thus treated is packed in an appropriate column  
15 to form the antibody column.

In purification with the above-mentioned antibody column, for example, a human BTC protein-containing sample is dissolved in a buffer having a neutral pH such as phosphate buffer or Tris-hydrochloric acid buffer, and  
20 adsorbed by the antibody column. Then, the column is washed with the same buffer, followed by elution of the human BTC protein. Eluents which can be used include slightly acidic solutions such as acetic acid solutions, solutions containing polyethylene glycol, solutions  
25 containing peptides more easily bindable with the antibody than the sample, high concentration salt solutions and combined solutions thereof. Solutions which do not so

- 26 -

accelerate decomposition of the human BTC protein are preferred.

Column eluates are neutralized with buffers by conventional methods. The above-mentioned purification  
5 procedure can be repeated if necessary.

Thus, the substantially pure human BTC protein substantially free from pyrogens and endotoxins is obtained. The substantially pure human BTC protein of the present invention contains the human BTC protein at a  
10 concentration of about 90% (w/w) or more, and preferably about 95% (w/w) or more.

The antibodies of the present invention bind to a human BTC protein or a mutein thereof with high sensitivity. Accordingly, they are very useful as reagents  
15 for assaying (detecting or determination) a human BTC protein or a mutein thereof or as reagents for purifying human BTC protein or a mutein thereof.

An abnormal expression of human BTC protein or a mutein thereof may cause diabetes or complications thereof.  
20 In these cases, the diseases can be foreseen by assaying the amount of human BTC protein. The sensitivity of the determination method is required to be as high as possible, because the amount of human BTC produced in vivo is very slight. According to the assays using the monoclonal  
25 antibodies of the present invention, the human BTC proteins can be measured down to 10 pg/ml, for example, as shown in Example 8 described below. This is very epochal in that in

- 27 -

vivo human BTC existing only in very slight amounts can be assayed. Diseases such as arterial sclerosis and acute tumors are considered as diseases induced by excess production of human BTC.

5       Accordingly, a diagnosis of diabetes or complications thereof can be conducted by using the antibody of the present invention. Diagnosis can be performed, for example, by the immunochemical assay of measuring the human BTC protein using an human BTC antibody bound to a carrier  
10 and a conjugate in which a labeling agent is directly bound to an anti-GAF antibody different from the former antibody in the antigen recognition site. The enzyme immunoassay is preferably employed among others.

      The antibodies of the present invention are used as  
15 both the antibody held on the carrier and the antibody different from the above-mentioned antibody in the antigen recognition site used in the above-mentioned assay, and the monoclonal antibodies are preferably used. Preferred examples of the monoclonal antibodies include antibodies  
20 1A1, 2B2 and 5E5 prepared in the example described below, and they may be appropriately selected.

      Carriers and labelling agents used in the detection and assay can also be used for the diagnosis. Examples of a diagnostic agent comprising the antibodies of the present  
25 invention include a combination of peroxidase-bound 5E5 antibody and 1A1 antibody bound to beads, and examples of a diagnosis kit include a combination of peroxidase-bound 5E5

antibody and 1A1 antibody bound to beads and a peroxidase substrate. By the diagnostic agent or the diagnosis kit, a human BTC protein can be assayed and accordingly diabetes or complications thereof be diagnosed.

5       As described above, the antibodies of the present invention have strong neutralizing activity, and therefore remove excess human BTC protein in vivo to neutralize activity of human BTC protein. For the diseases which might be induced by excess activity of human BTC protein,  
10   the antibodies can be given to humans as therapeutic agents.

The antibodies of the present invention are low in toxicity.

While the dosage of the antibodies of the present  
15   invention when administered to a patient varies with age, sex of the patient or state of disease, it ranges from about 100 µg/kg to about 10 mg/kg daily.

When the antibodies of the present invention are administered, the antibodies themselves or mixtures thereof  
20   with pharmaceutically acceptable carriers, excipients and diluents (for example, water and physiological saline) can be given parenterally in a formulation such as an injection.

Further, when parenteral preparations such as  
25   injections are prepared, isotonic agents (for example, glucose, D-sorbitol, D-mannitol and sodium chloride), preservatives (for example, benzyl alcohol, chlorobutanol,

methyl p-hydroxybenzoate and propyl p-hydroxybenzoate), buffers (for example, phosphate buffer and sodium acetate buffer), etc. can be appropriately added.

When nucleotides, amino acids and so on are indicated  
5 by abbreviations in the specification and drawings, the abbreviations adopted by the IUPAC-IUB Commission on Biochemical Nomenclature or commonly used in the art are employed. For example, the following abbreviations are used. When the amino acids are capable of existing as  
10 optical isomers, it is understood that the L-forms are represented unless otherwise specified.

	DNA	: Deoxyribonucleic acid
	A	: Adenine
	C	: Cytosine
15	G	: Guanine
	T	: Thymine
	Ala	: Alanine
	Arg	: Arginine
	Asn	: Asparagine
20	Asp	: Aspartic acid
	Cys	: Cysteine
	Gln	: Glutamine
	Glu	: Glutamic acid
	Gly	: Glycine
25	His	: Histidine
	Ile	: Isoleucine
	Leu	: Leucine

- 30 -

Lys : Lysine  
Met : Methionine  
Phe : Phenylalanine  
Pro : Proline  
5 Ser : Serine  
Thr : Threonine  
Trp : Tryptophan  
Tyr : Tyrosine  
Val : Valine

10 The present invention will be described in more detail through Reference Examples and Examples shown below. It is understood of course that they are not intended to limit the scope of the invention.

Transformant cell E. coli MM294(DE3)/pTB1516 obtained  
15 in Reference Example 3 described below was deposited with the National Institute of Bioscience and Human-technology (NIBH) [formerly the Fermentation Research Institute (FRI)], Agency of Industrial Science and Technology, the Ministry of International Trade and Industry, of 1-3,  
20 Higashi 1 chome, Tsukuba-shi Ibaraki-ken 305 Japan, under the accession number FERM BP-3836 on April 21, 1992, and with the Institute for Fermentation (IFO) of 17-85, Juso-honmachi 2-chome, Yodogawa-ku, Osaka 532 Japan, under the accession number IFO 15282 on April 16, 1992.

25 Transformant A9/1515-14 obtained in Reference Example 4 described below was deposited with the IFO of 17-85, Juso-honmachi 2-chome, Yodogawa-ku, Osaka 532 Japan, under



- 31 -

the accession number IFO 50389 on December 28, 1992, and with the NIBH of 1-3, Higashi 1 chome, Tsukuba-shi Ibaraki-ken 305 Japan under the accession number FERM BP-4159 on January 13, 1993.

5        Mouse 1A1 cell, 2B2 cell and 5E5 cell obtained in Example 2 described below were deposited with the IFO of 17-85, Juso-honmachi 2-chome, Yodogawa-ku, Osaka 532 Japan, under the accession numbers IFO 50437, 50438 and 50439 on February 15, 1994, and with the NIBH of 1-3, Higashi 1  
10 chome, Tsukuba-shi Ibaraki-ken 305 Japan under accession numbers FERM P-14847, FERM P-14848 and FERM P-14849 on March 23, 1995 which were transferred to international deposits under accession numbers FERM BP-5393, FERM BP-5394 and FERM BP-5395 on February 13, 1996.

15    Reference Example 1

Construction of Human BTC cDNA Expression Plasmid for Mammalian Cells

A 0.7-Kb SmaI-DraI fragment containing cDNA coding for human BTC protein (human BTC cDNA) was isolated from  
20 plasmid pTB1499 (refer to Example 4 of EP-A 0555785) [Biochem. Biophys. Res. Commun., 190, 1173 (1993)]. BglII linkers were ligated with this fragment at the flush ends thereof by use of T4 DNA ligase. After digestion with BglII, a 0.7-Kb fragment containing human BTC cDNA was  
25 inserted into the BglII site of expression plasmid pTB1308 prepared from pTB399 [Cell Structure Function, 12, 205 (1987)] by removing an IL-2 cDNA region, by ligation using

- 32 -

T4 DNA ligase (Fig. 1). Plasmid pTB1515 thus produced was cleaved with SalI and HindIII. A 2.4-Kb fragment containing MuLV LTR and human BTC cDNA was isolated and introduced between the SalI and HindIII sites of pTB348 [Cell Structure Function, 12, 205 (1987)] having an SV40 early region promoter and hamster DHFR cDNA. The resulting plasmid was named pTB1507 (Fig. 2).

#### Reference Example 2

Construction of Human BTC cDNA Expression Plasmid for  
10 E. coli

A 0.6-Kb EcoRI-BamHI fragment coding for a human BTC precursor (1-147 amino acid residues) was isolated from plasmid pTB1515 (Reference Example 1). A synthetic adaptor having an ATG translation initiation codon (a: 5' TATGGATGGG 3'; b: 5' AATTCCCATCCA 3') was ligated with the above-mentioned 0.6-Kb fragment at the EcoRI site thereof. Then, the resulting 0.6-Kb NdeI-BamHI fragment was inserted into plasmid pET-3c containing a T7 promoter [Gene, 56, 125 (1987)] to construct plasmid pTB1505.

20 In order to obtain a DNA fragment coding for 80 amino acid residues [from 1 (Asp) to 80 (Tyr) in Fig. 3] of human BTC protein, PCR was conducted using plasmid pTB1505 as a template and two oligonucleotides (1: 5' ATACATATGGATGGGAATTCCA 3' and 2: 5' CCGGATCCTAGTAAAACAAGTCAACTCT 3') as primers. The product was digested with NdeI and BamHI, and fractionated by 2.0% agarose gel electrophoresis, thereby isolating a desired

- 33 -

0.25-Kb DNA fragment. The 0.25-Kb NdeI-BamHI fragment was inserted downstream from the T7 promoter of pET-3c by ligation through T4 DNA ligase to obtain plasmid pTB1516 (Fig. 4).

### 5 Reference Example 3

#### Expression of Human BTC cDNA in E. coli

E. coli MM294 was lysogenized with lambda phage (study supra) recombined with an RNA polymerase gene of T7 phage. Then, plasmid pLysS was introduced into E. coli MM294(DE3) to obtain E. coli MM294(DE3)/pLysS. Plasmid pTB1516 was introduced into this strain, thereby obtaining E. coli MM294(DE3)/pLysS, pTB1516. The resulting transformant cells were cultivated in 20 ml of L-broth containing 100 µg/ml ampicillin and 10 µg/ml chloramphenicol at 37°C. When the Klett value was about 180, isopropyl-β-D-thiogalactoside (IPTG) was added to the medium to give a final concentration of 0.4 mM, and cultivation was continued for 4 hours. The resulting cells were collected, and suspended in 0.5 ml of a buffer containing 20 mM Tris-HCl (pH 7.4), 10 mM EDTA, 0.5 M NaCl, 10% sucrose and 0.25 mM PMSF, followed by addition of egg white lysozyme to this suspension at a concentration of 0.5 mg/ml. After placed in an ice bath for 1 hour, the mixed solution was cultivated at 37°C for 5 minutes, and centrifuged in a Sorvall centrifuge at 15,000 rpm at 4°C for 30 minutes to obtain a supernatant.

For this cell extract, DNA synthesis induction

- 34 -

activity was assayed by the uptake of  $^3\text{H}$ -thymidine into quiescent Balb/c 3T3 clone A31-714-4 cells [Int. J. Cancer, 12, 463 (1973)] according to the method described in Mol. Cell. Biol., 8, 588 (1988). Results thereof are shown in

5 Table 1.

In the measurement of human BTC protein biological activity, the DNA synthesis induction activity assay is usually used [T. Watanabe et al., J. Biol. Chem., 269 9966 (1994); EP-A 0555785].

10 Table 1

<u>Transduced DNA or control</u>	<u>Sample dilution</u>	<u><math>^3\text{H}</math>-thymidine uptake (cpm)</u>
<u>E. coli</u> MM294(DE3) /pLysS, pTB1516	1/78125 1/390625	20,232 13,169
<u>E. coli</u> MM294(DE3) /pLysS, pET:3c	1/3125	805 592

15

20

#### Reference Example 4

Establishment of human BTC-Producing A9 Cell Line

Mouse A9 cells (ATCC CCL 1.4) were co-transfected with

25 human BTC cDNA-containing plasmid pTB1515 (Reference Example 1) and human HPRT gene-containing expression plasmid p4aA8 [D. J. Jolly et al., Proc. Natl. Acad. Sci. USA, 80, 477 (1983)] by the calcium phosphate method. The resulting cells were proliferated in DMEM supplemented with

30 10% fetal calf serum for 2 days, followed by cultivation in HAT medium [J. W. Littlefield, Science, 145, 709 (1964)] for selection. HPRT<sup>+</sup> cells were proliferated in HAT medium, and clones were isolated by limiting dilution

- 35 -

method. Cells ( $10^5$  cells) of each clone were plated in each well of a 24-well plate, and cultivated in a growth medium for 2 days, followed by cultivation in 0.5 ml of DMEM containing 0.5% fetal calf serum for 2 days. The human BTC level at which  $10^6$  cells were secreted in the medium was examined by DNA synthesis induction activity on the mouse 3T3 cells described in Reference Example 3. Results of several clones are shown in Table 2.

Table 2

	<u>Clone</u>	<u>Activity (converted to mouse EGF, ng/ml)</u>
10	A9/1515-4	43
	A9/1515-14	566
	A9/1515-17	208
	A9/1515-34	258
15	A9/1515-63	94

Reference Example 5

## Purification of BTC Produced by A9 Cells

To 3.5 liters of a culture supernatant of the A9/1515-14 cells, 180 ml of 1 M potassium phosphate (pH 6.0), 7 ml of 0.5 M EDTA, 36 ml of 5% CHAPS and 7 ml of 0.25 M PMSF were added, and the resulting mixture was loaded on an S-Sepharose column (2.6 cm in diameter X 40 cm, Pharmacia). After the column was washed with 300 ml of a buffer [0.1 M potassium phosphate (pH 6.0), 1 mM EDTA, 0.05% CHAPS and 0.5 mM PMSF], the above-mentioned buffer containing 0.7 M NaCl was allowed to flow through the column at a flow rate

- 36 -

of 1 ml/minute to elute a protein. As to respective fractions collected for every 5 ml, the DNA synthesis induction activity (biological activity) of BALB/c3T3 cells was examined, and fraction Nos. 23-32 and 40-49 were pooled as BTC-I and BTC-II, respectively (Fig. 5).

The BTC-I pool fraction obtained from the S-Sepharose column was fractionated with 25% and 80% solutions of ammonium sulfate, followed by concentrating by ultrafiltration (Centriprep-10, Amicon). The concentrate was loaded at a flow rate of 1.2 ml/minute on a gel filtration column (1.6 cm in diameter X 60 cm; Superdex 75pg, Pharmacia) equilibrated with 20 mM Tris (pH 7.4), 1 mM EDTA and 0.05% CHAPS, and fractions were collected for every 1.2 ml from 15 minutes after the flowing initiation. A fraction which showed high biological activity (fraction No. 35-41) was pooled (Fig. 6A).

The BTC-II pool fraction obtained from the S-Sepharose column was concentrated by ultrafiltration (YM2, Amicon). The concentrate was loaded at a flow rate of 1.2 ml/minute on a gel filtration column (1.6 cm in diameter X 60 cm; Superdex 75pg, Pharmacia) equilibrated with 20 mM Tris (pH 7.4), 1 mM EDTA and 0.05% CHAPS, and fractions were collected from 15 minutes after initiation for every 1.2 ml. A fraction which showed high biological activity (fraction No. 66-74) was pooled (Fig. 6B).

The BTC-I pool fraction collected from the gel filtration column was loaded on a heparin HPLC column (0.8

- 37 -

in diameter X 5 cm; AFpak HR-894, Showa Denko K. K., Japan). The column was washed with 20 mM Tris-HCl (pH 7.4), 1 mM EDTA and 0.05% CHAPS, and then eluted with a linear gradient of 0-0.9 M NaCl at a flow rate of 1 ml/minute for 30 minutes to fractionate an eluate for every 1 ml. Fractions of fraction Nos. 9 to 13 in which biological activity was observed were pooled (Fig. 7A).

The BTC-II pool fraction collected from the gel filtration column was loaded on a heparin HPLC column (0.8 in diameter X 5 cm; AFpak HR-894, Showa Denko K. K., Japan). The column was washed with 20 mM Tris-HCl (pH 7.4), 1 mM EDTA and 0.05% CHAPS, and thereafter eluted with a linear gradient of 0-0.9 M NaCl at a flow rate of 1 ml/minute for 30 minutes to fractionate an eluate for every 1 ml. Fractions of fraction Nos. 16 to 19 in which biological activity was observed were pooled (Fig. 7B).

TFA was added to the BTC-I pool fraction collected from the heparin column to give a final concentration of 0.1%, and the TFA-containing fraction was loaded on a C18 reverse phase HPLC column (0.46 cm in diameter X 15 cm; Asahipak ODP-50, Asahi Chemical, Japan). Elution was performed with a linear gradient of 0-63% acetonitrile in the presence of 0.1% TFA for 70 minutes, and an eluate was collected for every 0.5 ml (1 minute) (Fig. 8A). Biological activity was observed in agreement with elution peaks of the protein. The protein in this portion was examined by SDS-PAGE/silver staining. As a result, a band

- 38 -

was detected only at the position corresponding to a molecular weight of 26 to 30 k (Fig. 9A).

By the above-mentioned operation, 150 µg of BTC-I was obtained.

5 TFA was added to the BTC-II pool fraction collected from the heparin column to give a final concentration of 0.1%, and the TFA-containing fraction was loaded on a C18 reverse phase HPLC column (0.46 cm in diameter X 15 cm; Asahipak ODP-50, Asahi Chemical, Japan). Elution was  
10 performed with a linear gradient of 0-63% acetonitrile in the presence of 0.1% TFA for 70 minutes, and an eluate was collected for every 0.5 ml (1 minute) (Fig. 8B).

Biological activity was observed in agreement with elution peaks of the protein. The protein in this portion was  
15 examined by SDS-PAGE/silver staining. As a result, a band was detected only at the position corresponding to a molecular weight of 14 k (Fig. 9B).

By the above-mentioned operation, 75 µg of BTC-II was obtained.

#### 20 Reference Example 6

Purification of BTC Produced from Recombinant E. coli  
E. coli MM294(DE3)/plysS, pTB1516 was cultivated overnight, and thereafter, LB medium was inoculated with the culture cell solution in a 20-fold dilution. After  
25 cultivation at 37°C for 2 hours, IPTG was added to give a final concentration of 0.1 mM, followed by further cultivation for 3 hours. Cells were collected by



- 39 -

centrifugation, and stored at -20°C until used.

The cell stock corresponding to the 5-liter culture was defrosted, and suspended in 300 ml of an ice-cooled buffer containing 50 mM Tris-HCl (pH 7.4), 10 mM EDTA, 0.2 M NaCl, 10% sucrose and 1 mM APMSF. Then, 40 mg of egg white lysozyme was dissolved therein. After incubation at 4°C for 2 hours, the suspension was subjected to ultrasonic treatment, and centrifuged at 20,000 X g for 1 hour to obtain a supernatant. This supernatant was allowed to pass through a 200-ml Q-Sepharose bed, and then TCA was added thereto to give a final concentration of 4%, followed by standing at 40°C for 10 minutes. A precipitate collected by centrifugation at 20,000 X g for 20 minutes was suspended in 100 ml of a buffer containing 20 mM Tris (pH 7.4), 1 mM EDTA, 0.15 M NaCl and 1 mM APMSF, and 5 M NaOH was added to the resulting suspension while homogenizing it with a moter to adjust the pH to 6. This homogenate was centrifuged at 100,000 X g for 1 hour to obtain a supernatant, which was loaded on an S-Sepharose column (1.6 cm in diameter X 10 cm, Pharmacia). The column was washed with a buffer containing 0.1 M potassium phosphate (pH 6.0), 1 mM EDTA and 0.5 mM PMSF, and thereafter eluted with 400 ml linear gradient of NaCl from 0 to 1 M for 200 minutes. An eluate was collected for every 5 ml. Fraction Nos. 20 to 27 and fraction Nos. 40 to 45 in which high biological activity was observed were pooled as E. coli BTC-I and E. coli BTC-II fractions, respectively (Fig. 10).

- 40 -

TFA was added to the BTC-I pool fraction to give a final concentration of 0.1%, and the TFA-containing fraction was loaded on a C18 reverse phase HPLC column (1.0 cm in diameter X 25 cm; Asahipak ODP-50, Asahi Chemical, Japan). The column was washed with 0.1% TFA, and thereafter eluted with 340 ml linear gradient of acetonitrile from 0% to 63% for 170 minutes. Then, an eluate was collected for every 4 ml. Biological activity was observed in agreement with the peak indicated by the arrow (Fig. 11). This peak was examined by SDS-PAGE/silver staining. As a result, a band was observed at the position corresponding to a molecular weight of about 18 k (Fig. 12, lane I). BTC-I was purified. By this method, 630 µg of E. coli BTC-I was obtained.

The BTC-I pool fraction was loaded on a C18 reverse phase HPLC column (4.6 cm in diameter X 15 cm; Asahipak ODP-50, Asahi Chemical, Japan) in the presence of 0.1% TFA. After washing, the column was eluted with 35 ml linear gradient of acetonitrile from 0% to 63% for 70 minutes, and an eluate was collected for every 0.5 ml. Biological activity was observed in agreement with the peak indicated by the arrow (Fig. 13). This peak was examined by SDS-PAGE/silver staining. As a result, a band was observed at the position corresponding to a molecular weight of less than 14.3 k (lysozyme) (Fig. 12, lane II). BTC-II was purified. By this method, 990 µg of E. coli BTC-II was obtained.

- 41 -

N-terminal amino acid sequences of E. coli BTC-I and E. coli BTC-II were determined up to 20 amino acid residues. The results showed that BTC-I had an N-terminal sequence starting from translation initiation methionine as expected, and that BTC-II had a molecule lacking 31 residues at the N-terminal side (Fig. 14).

#### Example 1

##### Immunization

Human BTC-I obtained in Reference Example 6 described above was dissolved in physiological saline, and the resulting solution was thoroughly suspended in an equivalent volume of Freund's complete adjuvant to prepare a suspension. The suspension was given intracutaneously to BALB/c mice (female, 10 weeks old) in a dose of 50 µg of antigen human BTC-I per mouse to perform an initial immunization. After 3 weeks, a suspension of human BTC-I in Freund's incomplete adjuvant was prepared and given to the mice in a manner similar to that of the initial immunization, thereby performing additional immunization. The booster was further conducted twice for every 2 weeks, and after 3 weeks, 50 µg of human BTC-I dissolved in physiological saline was given intraperitoneally to the mice to perform a final immunization.

#### Example 2

##### (1) Cell Fusion

Three days after the final immunization, the spleens were removed from the mice immunized in Example 1 to obtain

- 42 -

spleen cells to be used for cell fusion. These cells were suspended in RPMI 1640 medium (Nippon Seiyaku, Japan).

Mouse myeloma cells P3X63Ag 8UI (P3U1) were subcultured in RPMI 1640 medium supplemented with 10% fetal calf serum (FBS, Hyclon Laboratories). Cell fusion was carried out according to the method established by Köhler and Milstein [G. Köhler and C. Milstein, Nature, 256, 495 (1975)]. The resulting spleen cells and the mouse myeloma cells were mixed in a ratio of 5:1. After sufficient removal of the medium, the mixture was incubated at 37°C for 2 minutes in 50% polyethylene glycol 4000 (Sigma) dissolved in 1 ml of RPMI 1640 medium. The resulting cells were washed with serum-free RPMI 1640 medium, and then suspended in RPMI-1640 medium (containing 10% FBS) supplemented with HAT ( $1 \times 10^{-4}$  M hypoxanthine,  $4 \times 10^{-7}$  M aminopterin and  $1.6 \times 10^{-5}$  M thymidine) (hereinafter referred to as HAT medium). The suspension was dispensed into each well of a 96-well culture plate by each  $3 \times 10^5$  cells/0.1 ml. After cultivation at 37°C in 7% CO<sub>2</sub> for 3 days, 0.1 ml of HAT medium was further added to each well. The cells which grew under these conditions were hybridomas.

## (2) Screening for Antibody-Producing Cells

Physiological saline containing 10 µg/ml of BTC-I obtained in Reference Example 5 was added in an amount of 0.1 ml to each well of a 96-well flat-bottomed ELISA plate (Falcon 3912), followed by standing at 4°C overnight. After washing three times with 0.05% Tween 20-PBS, 0.2 ml

- 43 -

of 5 mg/ml BSA-0.05% Tween 20-PBS was added to each well, followed by standing at 37°C for 1 hour. A culture supernatant which was subjected to a serial 2-fold stepwise dilution with 0.05% Tween 20-PBS was added in an amount of 0.1 ml/well, and the plate was allowed to stand at room temperature for 2 hours. After washing three times with 0.05% Tween 20-PBS, 0.1 ml of an HRP-labeled anti-mouse IgG rabbit antibody solution (1 µg/ml) was added to each well as a secondary antibody, followed by standing at room temperature for 1 hour. After washing was repeated 5 times, 0.1 ml of a color-developing solution (prepared by mixing 10 ml of o-phenylenediamine, 10 ml of 50 mM disodium phosphate-24 mM citrate buffer and 120 µl of a 1.7% hydrogen peroxide solution at the time of use) was added to each well to conduct HRP enzyme reaction. The enzyme reaction was terminated by addition of 0.05 ml of 6 N sulfuric acid, and the absorbance at a wavelength of 492 nm was measured (ELISA).

By this assay, primary screening was performed for culture supernatants of wells in which the hybrid cells grew, and cells in wells which exhibited high absorbance were cloned by the limiting dilution analysis. In cloning, the cells were preliminarily cultivated at 37°C in 7% CO<sub>2</sub> overnight. Mouse peritoneal exudate cells (10<sup>4</sup> cells/well) were used as a feeder. Screening and cloning were further repeated to obtain anti-human BTC antibody-producing hybridomas 1A1, 2B2 and 5E5. When the antibody titer was

- 44 -

assayed for hybridoma culture supernatants by the above-mentioned ELISA, the absorbance at a wavelength of 492 nm was 1 or more even when 128-fold dilutions were made. The cloned hybridoma cells were suspended in RPMI-1640 medium supplemented with 20% fetal calf serum and 10% dimethyl sulfoxide (DMSO) and stored in liquid nitrogen.

### Example 3

#### Immunoglobulin Class of Monoclonal Antibodies

Culture supernatants of the hybridoma cells obtained in Example 2-(2) were tested by a mouse antibody subclass detecting kit (Bio RAD) to determine immunoglobulin subclass. Results are shown in Table 3.

Table 3

15	Immunoglobulin subclass	<u>Hybridoma culture supernatant</u>		
		<u>1A1</u>	<u>2B2</u>	<u>5E5</u>
20	IgG1	+	+	+
	IgG2a	-	-	-
	IgG2b	-	-	-
25	IgG3	-	-	-
	IgM	-	-	-
	IgA	-	-	-
30				

In Table 3, "+" indicates the positive reaction, and "-" indicates the negative reaction.

The results shown in Table 3 reveal that three kinds of antibodies in the hybridoma culture supernatants belong to IgG1.

Example 4

## Recognition Site of Monoclonal Antibody

To each well of a 96-well flat-bottomed ELISA plate (Falcon 3912), 0.1 ml of physiological saline containing 1  
5  $\mu\text{g/ml}$  of BTC-I or BTC-II obtained in Reference Example 5 was added, followed by standing at  $4^{\circ}\text{C}$  overnight. After three time washings with 0.05% Tween 20-PBS, 0.2 ml of 5 mg/ml BSA-0.05% Tween 20-PBS was added to each well, followed by standing at  $37^{\circ}\text{C}$  for 1 hour. After three time  
10 washings with 0.05% Tween 20-PBS again, a culture supernatant which was subjected to a serial 2-fold stepwise dilution with 0.05% Tween 20-PBS was added in an amount of 0.1 ml/well, and the plate was allowed to stand at room temperature for 2 hours. After three time washings with  
15 0.05% Tween 20-PBS, 0.1 ml of an HRP-labeled anti-mouse IgG rabbit antibody solution ( $1 \mu\text{g/ml}$ ) was added to each well as a secondary antibody, followed by standing at room temperature for 1 hour. After washing was repeated 5 times, 0.1 ml of a color-developing solution (prepared by  
20 mixing 10 ml of o-phenylenediamine, 10 ml of 50 mM disodium phosphate-24 mM citrate buffer and 120  $\mu\text{l}$  of a 1.7% hydrogen peroxide solution at the time of use) was added to each well to conduct HRP enzyme reaction. The enzyme reaction was terminated by addition of 0.05 ml of 6 N  
25 sulfuric acid, and the absorbance at a wavelength of 492 nm was measured (ELISA). Results are shown in Table 4.

- 46 -

Table 4Hybridoma culture supernatant

	<u>1A1</u>	<u>2B2</u>	<u>5E5</u>
Human BTC-I	+	+	+
5 Human BTC-II	+	+	+

In Table 4, "+" indicates the positive reaction, and "-" indicates the negative reaction.

The results shown in Table 4 reveal that three kinds of antibodies in the hybridoma culture supernatants recognize the C-terminal side from Arg<sup>62</sup> of human BTC (+31 in Fig. 3).

Example 5

## Reaction Specificity of Monoclonal Antibody

To each well of a 96-well flat-bottomed ELISA plate (Falcon 3912), 0.1 ml of physiological saline containing 1 µg/ml of BTC-I obtained in Reference Example 5, Human TGF-α (Gibco-BRL), human EGF (Boehringer Mannheim) or mouse EGF (Takara Shuzo, Japan) was added, followed by standing at 4°C overnight. After three time washings with 0.05% Tween 20-PBS, 0.2 ml of 5 mg/ml BSA-0.05% Tween 20-PBS was added to each well, followed by standing at 37°C for 1 hour. After three time washings with 0.05% Tween 20-PBS again, a culture supernatant which was subjected to a serial 2-fold stepwise dilution with 0.05% Tween 20-PBS was added in an amount of 0.1 ml/well, and the plate was allowed to stand at room temperature for 2 hours. After three time washings with 0.05% Tween 20-PBS, 0.1 ml of an HRP-labeled anti-



- 47 -

mouse IgG rabbit antibody solution (1 µg/ml) was added to each well as a secondary antibody, followed by standing at room temperature for 1 hour. After five time washings were repeated, 0.1 ml of a color-developing solution (prepared by mixing 10 ml of o-phenylenediamine, 10 ml of 50 mM disodium phosphate-24 mM citrate buffer and 120 µl of a 1.7% hydrogen peroxide solution at the time of use) was added to each well to conduct HRP enzyme reaction. The enzyme reaction was terminated by addition of 0.05 ml of 6 N sulfuric acid, and the absorbance at a wavelength of 492 nm was measured (ELISA). Results are shown in Table 5.

Table 5Hybridoma culture supernatant

		<u>1A1</u>	<u>2B2</u>	<u>5E5</u>
15	BTC-I	+	+	+
	Human TGF-α	-	-	-
	Human EGF	-	-	-
	Mouse EGF	-	-	-

In Table 5, "+" indicates the positive reaction, and  
20 "-" indicates the negative reaction.

The results shown in Table 5 reveal that three kinds of antibodies in the hybridoma culture supernatants specifically recognize BTC.

Example 6

25       Purification of Monoclonal Antibody from Culture Supernatant

A 1:1 mixture of the culture supernatant of mouse

- 48 -

hybridoma 1A1, 2B2 or 5E5 obtained in Example 2 and a binding buffer [3 M sodium chloride and 1.5 M glycine (pH 8.7)] was loaded onto a protein A column equilibrated with the binding buffer to allow an antibody to be adsorbed thereby. After washing with the binding buffer, the antibody was eluted with an eluting buffer [0.1 M citric acid (pH 5)]. 1 M Tris (pH 8.0) was added to the eluate to neutralize it, followed by dialysis with phosphate buffered saline. Thus, purified monoclonal antibody 1A1, 2B2 or 5E5 was obtained.

#### Example 7

##### Assay of Human BTC-Neutralizing Activity

For antibodies 1A1, 2B2 and 5E5 purified in Example 6, neutralizing activity on human BTC was assayed. Mouse Balb/c 3T3 clone A31-714-4 cells were seeded on a 96-well plate at  $2 \times 10^3$  cells per well, and cultivated in 100  $\mu$ l of DMEM supplemented with 5% fetal calf serum. The next day, the medium was changed by DMEM supplemented with 1% FCS, and cultivation was further continued for 3 days. Then, 1 ng/ml of purified human BTC-I, as well as 0.1, 1 and 10  $\mu$ g/ml of the above-mentioned three kinds of monoclonal antibodies to human BTC, was added thereto. After cultivation for 18 hours, 0.5  $\mu$ Ci/well of  $^3$ H thymidine (5 Ci/mmol) was added, followed by cultivation for 6 hours. The culture solution was removed, and the cells were washed 3 times with PBS. Then, 100  $\mu$ l of 5% SDS was added to dissolve the cells. The radioactivity taken into cell DNAs

was measured by use of a liquid scintillation counter to assay the neutralizing activity of the respective monoclonal antibodies on the DNA synthesis induction ability due to human BTC. Results are shown in Fig. 15.

5 Referring to Fig. 15, ■, □, ◇ and △ indicate the uptake values of <sup>3</sup>H-thymidine when no antibody, antibody 1A1, antibody 2B2 and antibody 5E5 were added, respectively. All the monoclonal antibodies inhibited DNA synthesis of the cells when human BTC was added, depending  
10 on the concentration. In particular, antibody 5E5 exhibited high neutralizing activity.

On the other hand, similar experiments were conducted for human EGF, human TGF- $\alpha$ , human HB-EGF and mouse BTC belonging to the same EGF family. The DNA synthesis of A31  
15 cells induced by addition of 1 ng/ml of each growth factor was not inhibited at all even when 10  $\mu$ g/ml of antibody 1A1, 2B2 or 5E5.

The results described above proved that all of monoclonal antibodies 1A1, 2B2 and 5E5 specifically  
20 neutralized the biological activity of human BTC protein.

#### Example 8

(1) One milligram of monoclonal antibody 5E5 obtained in Example 6 was dialyzed against 0.1 M NaHCO<sub>3</sub> (pH 8.0). Then, 120  $\mu$ g of NHS-biotin (Vector, U.S.A.) dissolved in  
25 DMSO was added to the dialyzed antibody, followed by reaction at room temperature for 2 hours with stirring. After reaction, the reaction product was dialyzed against

- 50 -

PBS(-) to remove unreacted NHS-biotin.

(2) Monoclonal antibody 1A1 obtained in Example 6 was diluted with 0.01 M  $\text{Na}_2\text{HPO}_4$  (pH 8.0) supplemented with 0.01 M NaCl to a concentration of 10  $\mu\text{g/ml}$ . The resulting  
5 solution was poured in an amount of 50  $\mu\text{l/well}$  into an immunoplate (Nunc, Denmark), followed by standing overnight at 4°C to allow it to be adsorbed by the plate. Antibodies not adsorbed were removed, followed by three time washings with PBS(-). A blocking solution [a solution prepared by  
10 diluting Block Ace (Dainippon Seiyaku, Japan) four times with PBS(-) and containing 0.01% Mlthiolate (trade mark)] in an amount of 100  $\mu\text{l/well}$ . After standing at 4°C overnight, the blocking solution was removed, followed by five time washings with PBS(-). Then, human BTC protein  
15 diluted with a diluent [a solution prepared by adding 0.2% Tween 20 to PBS(-)] was added in an amount of 50  $\mu\text{l/well}$ , followed by reaction at 37°C for 2 hours. After removal of unreacted BTC, the resulting product was washed 6 times with a washing solution [a solution prepared by adding  
20 0.05% Tween 20 to PBS(-)]. Further, biotinated monoclonal antibody 5E5 prepared in the above (1) was diluted 1:2000 with the diluent, and the diluted product was added in an amount of 50  $\mu\text{l/well}$ , followed by reaction at 37°C for 2 hours. After removal of an antibody-containing solution,  
25 the resulting product was washed 6 times with the washing solution, and avidin D-binding horseradish peroxidase (Vector, U.S.A.) diluted 1:1000 with the diluent was added

in an amount of 50  $\mu$ l/well, followed by reaction at room temperature for 30 minutes. After removal of the reaction solution, the resulting product was washed 8 times with the washing solution, and 50  $\mu$ l/well of peroxidase substrate  
5 (Wako, Japan) was added to develop color. Thereafter, 50  $\mu$ l/well of 2 N sulfuric acid was added to terminate reaction. After termination of the reaction, the colorimetric assay was performed.

Fig. 16 shows a detection curve of human BTC protein.  
10 The numbers on the abscissa indicate the concentration of human BTC protein added, and the numbers on the ordinate indicate the absorbance (at A492 nm) measured. The absorbance when BTC protein was not added was 0.06 to 0.07. When the concentration of human BTC showing an at least  
15 twice absorbance that was regarded as significant, the above-mentioned assay system was able to detect and determine human BTC having a concentration of 10 to 200 pg/ml.

In this assay system, human EGF, mouse EGF, human TGF-  
20  $\alpha$ , human HB-EGF and mouse BTC only showed background levels of absorbances, as shown in Table 6, which proved that this assay system was specific for human BTC protein.

- 52 -

Table 6

<u>Specimen</u>	<u>Concentration (ng/ml)</u>	<u>Absorbance (A<sub>492</sub>)</u>
-	0	0.062
Human EGF	1000	0.069
5 Mouse EGF	1000	0.064
Human TGF- $\alpha$	1000	0.077
Human HB-EGF	1000	0.066
Mouse BTC	300	0.069
Human BTC-I	100	4.063
10 Human BTC-I	0.1	0.889

Example 9

## Purification of Human BTC Using Antibody Column

Twenty milligrams of monoclonal antibody 5E5 obtained in Example 6 was allowed to bind to 10 g of a wet gel of Formyl-Cellulofine (Seikagaku Corporation, Japan), and loaded onto an econo column to prepare an affinity column.

On the other hand, plasmid pTB1685 (a derivative of plasmid pTB1507 described in Reference Example 1; a plasmid in which the MuLV-LTR region was changed to the human EF-1 promoter (Nuclei Acids Res., 18, 5322), and the human BTC cDNA region was changed to the cDNA region coding for BTC-I having the amino acid sequence represented by SEQ ID NO: 1, using plasmid pTB1516 described in Reference Example 2 as a material) was introduced into hamster CHO DHFR<sup>-</sup> cells, and human BTC high producing cell CHO 1685-39 was obtained from the cell lines converted to DHFR<sup>+</sup>. After the CHO 1685-39 cells were cultivated to a confluent in DMEM supplemented

- 53 -

with 5% fetal calf serum and 35 µg /ml proline, the medium was changed by ASF medium (Ajinomoto, Japan), and the cells were further cultivated for 3 days, followed by recovery of a culture supernatant. Five liters of the culture supernatant was concentrated about 4-fold on an Amicon concentrator under ice cooling, and then the resulting concentrate was allowed to flow through the above-mentioned affinity column at a flow rate of 20 ml/hour to allow BTC to be adsorbed thereby. After adsorption, the column was washed with Dulbecco's phosphate buffer (PBS(-)) until the absorbance at 280 nm reached 0.05 or less. Then, the column was washed with 0.01 M phosphate buffer (pH 6.5) containing 0.5 M NaCl at the same flow rate. Further, the column was eluted with 0.05 M glycine-HCl (pH 2.3) containing 0.15 M NaCl at the same flow rate. The eluted fractions were neutralized with 1 M Tris-HCl (pH 8.0), and dialyzed against PBS(-).

The resulting fractions were subjected to 15% acrylamide electrophoresis [Laemmli, Nature, 277, 680 (1970)]. The Coomassie Blue staining method was employed for staining.

Fig. 17 shows results for (1) a molecular weight marker, (2) the culture supernatant, (3) a fraction not adsorbed to the column, (4) a fraction eluted at a high salt-concentration (0.5 M NaCl) and (5) a fraction eluted with an eluting buffer.

Further, trifluoroacetic acid (TFA) was added to the

- 54 -

eluted fraction (5) to give a final concentration of 0.05%, and the TFA-containing fraction was loaded on a TSK gel-ODS column (Tosoh, Japan). Elution was performed with a linear gradient of 0-60% acetonitrile in the presence of 0.05% TFA for 60 minutes, and an eluate was collected at 1 ml/minute (Fig. 18). Elution peaks a, b and c of the protein were subjected to 15% acrylamide electrophoresis, and stained by the silver staining method (Fig. 19). Numerals (1), (5), (6), (7) and (8) indicate the molecular weight marker, the fraction eluted from the affinity column, elution peak a, elution peak b and elution peak c, respectively.

For each fraction obtained in the above-mentioned purifying course, the amount of huma BTC protein was assayed by the method described in Example 8. Results are shown in Table 7.



- 55 -

Table 7

	<u>Eluted fractions</u>	<u>Total amount of protein (A) (<math>\mu</math>g)</u>	<u>Amount of BTC (B) (<math>\mu</math>g)</u>	<u>(B)/(A)</u>
5	Culture supernatant	5360000	637.0	0.00012
	Concentrated culture supernatant (2)	1582600	563.2	0.00036
	Fraction not adsorbed by affinity column (3)	1267400	25.9	0.00002
10	Fraction eluted with phosphate buffer (pH 6.5) + 0.5 M NaCl (4)	336	1.9	0.00565
	Fraction eluted with 0.01 M glycine-HCl (pH 2.3) + 0.5 M NaCl (5)	750	371.9	0.50
15	Reverse phase HPLC Peak a (6)	88	66.9	0.76
	Peak b (7)	158	131.4	0.83
20	Peak c (8)	124	130.5	1.05

The biological activity of the above-mentioned eluted fractions was assayed by the uptake experiment of  $^3\text{H}$ -thymidine using the mouse Balb/c 3T3 cells described in Example 7. As a result, all the fractions showed DNA synthesis stimulating activity corresponding to the amount of human BTC protein.

The above results revealed that the use of the antibody columns using the antibody of the present invention allowed efficient purification of human BTC protein.

Example 10

Preparation of Anti-Peptide Antibody (polyclonal

- 56 -

antibody)

A polypeptide having the amino acid sequence from the 66th to the 80th of the amino acid sequence represented by SEQ ID NO: 1 was chemically synthesized to use it as an  
5 antigen.

(1) The above-mentioned peptide (5 mg) and hemocyanin (10 mg) were dissolved in 4 ml of 0.2 M phosphate buffer (pH 7.3), and 400  $\mu$ l of 2.5% glutaraldehyde cooled in ice water was added thereto drop by drop with stirring. After  
10 stirring for 3 hours under ice cooling, the solution was dialyzed against distilled water to obtain a conjugate of the peptide and hemocyanin.

(2) Bovine serum albumin (BSA) (132 mg) was dissolved in 3 ml of 0.1 M phosphate buffer (pH 7.5) to obtain  
15 solution A. On the other hand, 11.2 mg of N-( $\gamma$ -maleimidobutyloxy)succinimide (GMBS) was dissolved in 200  $\mu$ l of a dimethylformamide solution to obtain solution B. Solution B was added dropwise to solution A while stirring with a stirrer, and the mixed solution was reacted at 30°C  
20 for 30 minutes. Then, the resulting solution was purified on a Sephadex-G-25 column (1.5 cm X 20 cm) using 0.1 M phosphate buffer (pH 6.5)-0.1 M NaCl as an eluent to obtain bovine serum albumin into which a maleimido group was introduced.

25 The peptide (5 mg) was dissolved in 0.1 M phosphate buffer-5 mM ethylenediaminetetraacetic acid (EDTA), and maleimido group-introduced bovine serum albumin (20 mg) was

- 57 -

added thereto (to a total amount of 5 ml or less), followed by reaction at 30°C for 60 minutes. PBS was added thereto to bring the volume to 12 ml, thereby obtaining a conjugate of the peptide and bovine serum albumin.

5       (3) The conjugate of the peptide and hemocyanin obtained in the above (1) was thoroughly mixed with Freund's complete adjuvant, and rabbits were subcutaneously injected with the resulting mixture. Thereafter, the conjugate of the peptide and bovine serum albumin obtained  
10 in the above (2) was mixed with Freund's complete adjuvant, and the same rabbits were injected with the resulting mixture every two weeks. The blood collected from the rabbits immunized by the above-mentioned method was centrifuged to obtain a human BTC peptide antibody  
15 (polyclonal antibody).

The antibodies of the present invention neutralize biological activity of the human BTC protein or a mutein thereof, and bind to the protein or a mutein thereof with high sensitivity and high specificity. They can be  
20 therefore used as therapeutic agents for diseases such as arterial sclerosis and cancers. They can be also used as reagents for detecting and assaying human BTC protein or a mutein thereof, and also used as a diagnostic agent for diabetes or complications thereof.

- 58 -

## SEQUENCE LISTING

## (2) INFORMATION FOR SEQ ID NO:1:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 80 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: protein

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

Asp	Gly	Asn	Ser	Thr	Arg	Ser	Pro	Glu	Thr	Asn	Gly	Leu	Leu	Cys	Gly	1	5	10	15
Asp	Pro	Glu	Glu	Asn	Cys	Ala	Ala	Thr	Thr	Thr	Gln	Ser	Lys	Arg	Lys	20	25	30	
Gly	His	Phe	Ser	Arg	Cys	Pro	Lys	Gln	Tyr	Lys	His	Tyr	Cys	Ile	Lys	35	40	45	
Gly	Arg	Cys	Arg	Phe	Val	Val	Ala	Glu	Gln	Thr	Pro	Ser	Cys	Val	Cys	50	55	60	
Asp	Glu	Gly	Tyr	Ile	Gly	Ala	Arg	Cys	Glu	Arg	Val	Asp	Leu	Phe	Tyr	65	70	75	80

## (2) INFORMATION FOR SEQ ID NO:2:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1271 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: DNA (genomic)

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo sapiens
- (B) STRAIN: MCF7
- (C) INDIVIDUAL ISOLATE: Breast Adenocarcinoma Cell

## (ix) FEATURE:

- (A) NAME/KEY: CDS
- (B) LOCATION: 295..828

## (ix) FEATURE:

- (A) NAME/KEY: mat\_peptide
- (B) LOCATION: 388..828

- 59 -

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

CAGCGTGGAG GCTCCAAGGA CCAAGTCCTG CGCCTCTTTG GCGGGGTGTG TGCAGGAGGA																60
GGGGGGATAA ATAGGAGGCT CCCTCCTCCC GGCGACATTC ACGGAGCCGG CCGGCCTCCC																120
GCCCTGGGTG TTTCCCTGCC TTGTAGCCAG GGTGCCAGCC TGGGAAGTAG TTTCGTTTCC																180
TTCTGCCTCC GGGATTAGTT TCCAGGCACC CTCTCAGGCG CCCGAGGCCG GGAAGGGGG																240
CGAAGAAGGA GGGAGACTTG TCTAGGGGCT GCCCGGCCCG GCAGAGCGGG GTTG ATG																297
																Met
																-31
GAC CGG GCC GCC CGG TGC AGC GGC GCC AGC TCC CTG CCA CTG CTC CTG																345
Asp Arg Ala Ala Arg Cys Ser Gly Ala Ser Ser Leu Pro Leu Leu Leu																-15
-30 -25 -20																
GCC CTT GCC CTG GGT CTA GTG ATC CTT CAC TGT GTG GTG GCA GAT GGG																393
Ala Leu Ala Leu Gly Leu Val Ile Leu His Cys Val Val Ala Asp Gly																1
-10 -5																
AAT TCC ACC AGA AGT CCT GAA ACT AAT GGC CTC CTC TGT GGA GAC CCT																441
Asn Ser Thr Arg Ser Pro Glu Thr Asn Gly Leu Leu Cys Gly Asp Pro																
5 10 15																
GAG GAA AAC TGT GCA GCT ACC ACC ACA CAA TCA AAG CGG AAA GGC CAC																489
Glu Glu Asn Cys Ala Ala Thr Thr Thr Gln Ser Lys Arg Lys Gly His																
20 25 30																
TTC TCT AGG TGC CCC AAG CAA TAC AAG CAT TAC TGC ATC AAA GGG AGA																537
Phe Ser Arg Cys Pro Lys Gln Tyr Lys His Tyr Cys Ile Lys Gly Arg																50
35 40 45																
TGC CGC TTC GTG GTG GCC GAG CAG ACG CCC TCC TGT GTC TGT GAT GAA																585
Cys Arg Phe Val Val Ala Glu Gln Thr Pro Ser Cys Val Cys Asp Glu																65
55 60																
GGC TAC ATT GGA GCA AGG TGT GAG AGA GTT GAC TTG TTT TAC CTA AGA																633
Gly Tyr Ile Gly Ala Arg Cys Glu Arg Val Asp Leu Phe Tyr Leu Arg																70 75 80
70 75 80																
GGA GAC AGA GGA CAG ATT CTG GTG ATT TGT TTG ATA GCA CTT ATG GTA																681
Gly Asp Arg Gly Gln Ile Leu Val Ile Cys Leu Ile Ala Val Met Val																
85 90 95																
GTT TTT ATT ATT TTG GTC ATC GGT GTC TGC ACA TGC TGT CAC CCT CTT																729
Val Phe Ile Ile Leu Val Ile Gly Val Cys Thr Cys Cys His Pro Leu																
100 105 110																
CGG AAA CGT CGT AAA AGA AAG AAG AAA GAA GAA GAA ATG GAA ACT CTG																777
Arg Lys Arg Arg Lys Arg Lys Lys Lys Glu Glu Glu Met Glu Thr Leu																115 120 125 130

- 60 -

GGT AAA GAT ATA ACT CCT ATC AAT GAA GAT ATT GAA GAG ACA AAT ATT 825  
 Gly Lys Asp Ile Thr Pro Ile Asn Glu Asp Ile Glu Glu Thr Asn Ile  
                   135                                  140                                  145

GCT TAAAAGGCTA TGAAGTTACC TCCAGGTTGG TGGCAAGCTG CAAAGTGCCT 878  
 Ala

TGCTCATTTG AAAATGGACA GAATGTGTCT CAGGAAAAAC AGCTAGTAGA CATGAATTTT 938  
 AAATAATGTA TTTACTTTTT ATTTGCAACT TTAGTTTGTG TTATTATTTT TTAATAAGAA 998  
 CATTAATTAT ATGTATATTG TCTAGTAATT GGGAAAAAAG CAACTGGTTA GGTAGCAACA 1058  
 ACAGAAGGGA AATTTCAATA ACCTTTCCTT TAAGTATTGT CACCAGGATT ACTAGTCAAA 1118  
 CAAAAAAGAA AAGTAGAAAG GAGGTTAGGT CTTAGGAATT GAATTAATAA TAAAGCTACC 1178  
 ATTTATCAAG CATTTACCAT GTGCTAATAA GTTTGAAATA TATTATTTCC TTTATTCCTT 1238  
 TCAGCAATCC ATGAGATAGC TATTATAATC CTC 1271

(2) INFORMATION FOR SEQ ID NO:3:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 80 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

Asp	Gly	Asn	Thr	Thr	Arg	Thr	Pro	Glu	Thr	Asn	Gly	Ser	Leu	Cys	Gly
1				5					10					15	
Ala	Pro	Gly	Glu	Asn	Cys	Thr	Gly	Thr	Thr	Pro	Arg	Gln	Lys	Val	Lys
			20					25					30		
Thr	His	Phe	Ser	Arg	Cys	Pro	Lys	Gln	Tyr	Lys	His	Tyr	Cys	Ile	His
		35					40					45			
Gly	Arg	Cys	Arg	Phe	Val	Val	Asp	Glu	Gln	Thr	Pro	Ser	Cys	Ile	Cys
	50					55					60				
Glu	Lys	Gly	Tyr	Phe	Gly	Ala	Arg	Cys	Glu	Arg	Val	Asp	Leu	Phe	Tyr
65					70				75					80	

## WHAT IS CLAIMED IS:

1. An antibody which specifically binds to human betacellulin protein or a mutein thereof.
2. The antibody according to claim 1 wherein human betacellulin protein has an amino acid sequence of SEQ ID NO:1.
3. The antibody according to claim 1 or 2 which is a monoclonal antibody.
4. The antibody according to anyone of claims 1 to 3 which has no cross-reactivity with at least one of human epidermal growth factor, human transforming growth factor  $\alpha$  and mouse betacellulin protein.
5. A monoclonal antibody which specifically binds to human betacellulin protein and is capable of neutralizing the biological activity of said protein, which does not have cross-reactivity with human epidermal growth factor and human transforming growth factor  $\alpha$ , and which belongs to the immunoglobulin class of IgG.
6. The antibody according to anyone of claims 1 to 5 which recognizes an amino acid sequence of 31st (Arg) to 80th (Tyr) of SEQ ID NO:1.
7. A cloned hybridoma which is composed of a mammalian spleen cell immunized by human betacellulin protein or a mutein thereof and a homogenic or heterogenic lymphocyte.
8. A method of producing the monoclonal antibody according to claim 5, which comprises proliferating the hybridoma according to claim 7 in a liquid culture medium or

in an abdominal cavity of a mammal to form and accumulate the monoclonal antibody and then collecting the monoclonal antibody.

9. A method of detecting and assaying a human betacellulin protein or a mutein thereof which comprises contacting the antibody according to anyone of claims 1 to 6 with a specimen.

10. A method of diagnosing diabetes or complications thereof which comprises contacting the antibody according to anyone of claims 1 to 6 with a specimen and assaying a human betacellulin protein or a mutein thereof in the specimen.

11. A pharmaceutical composition which comprises an effective amount of the antibody according to anyone of claims 1 to 6 and a pharmaceutically acceptable carrier, excipient or diluent.

12. The pharmaceutical composition according to claim 11 for diagnosing diabetes or complications thereof.

13. A kit for diagnosing diabetes or complications thereof which comprises an effective amount of the antibody according to anyone of claims 1 to 6.


14. A kit for detecting and assaying a human betacellulin protein or a mutein thereof which comprises an effective amount of the antibody according to anyone of claims 1 to 6.

15. A method of purifying a human betacellulin protein or a mutein thereof which comprises contacting the antibody according to anyone of claims 1 to 6 with a crude sample containing the protein or a mutein thereof and isolating it.




1 / 20

FIG.1



2 / 20

FIG. 2



3 / 2 0

F I G . 3 - 1

10 20 30 40 50 60  
 CAGCGTGGAGGCTCCAAGGACCAAGTCCTGCGCCTCTTTGGCGGGGTGTGTGCAGGAGGA  
 70 80 90 100 110 120  
 GGGGGGATAAATAGGAGGCTCCCTCCTCCCGGCGACATTACGGAGCCGGCCGGCCTCCC  
 130 140 150 160 170 180  
 GCCCTGGGTGTTTCCCTGCCTTGTAGCCAGGGTGCCAGCCTGGGAAGTAGTTTCGTTTCC  
 190 200 210 220 230 240  
 TTCTGCCTCCGGGATTAGTTTCCAGGCACCCTCTCAGGCGCCCGAGGCCCCGGGAAGGGGG  
 250 260 270 280 290 300  
 CGAAGAAGGAGGGAGACTTGTCTAGGGGCTGCCCGGCCCGGCAGAGCGGGGTGATGGAC  
 Met Asp  
 -31  
 310 320 330 340 350 360  
 CGGGCCGCCCGGTGCAGCGGCGCCAGCTCCCTGCCACTGCTCCTGGCCCTTGCCCTGGGT  
 ArgAlaAlaArgCysSerGlyAlaSerSerLeuProLeuLeuLeuAlaLeuAlaLeuGly  
 370 380 390 400 410 420  
 CTAGTGATCCTTCACTGTGTGGTGGCAGATGGGAATTCCACCAGAAGTCCTGAACTAAT  
 LeuValIleLeuHisCysValValAlaAspGlyAsnSerThrArgSerProGluThrAsn  
 ↑ +1 10  
 430 440 450 460 470 480  
 GGCTCCTCTGTGGAGACCCTGAGGAAAACCTGTGCAGCTACCACCACACAATCAAAGCGG  
 GlyLeuLeuCysGlyAspProGluGluAsnCysAlaAlaThrThrThrGlnSerLysArg  
 20 30  
 490 500 510 520 530 540  
 AAAGGCCACTTCTCTAGGTGCCCCAAGCAATACAAGCATTACTGCATCAAAGGGAGATGC  
 LysGlyHisPheSerArgCysProLysGlnTyrLysHisTyrCysIleLysGlyArgCys  
 40 50  
 550 560 570 580 590 600  
 CGCTTCGTGGTGGCCGAGCAGACGCCCTCCTGTGTCTGTGATGAAGGCTACATTGGAGCA  
 ArgPheValValAlaGluGlnThrProSerCysValCysAspGluGlyTyrIleGlyAla  
 60 70  
 610 620 630 640 650 660  
 AGGTGTGAGAGAGTTGACTTGTTTTACCTAAGAGGAGACAGAGGACAGATTCTGGTGATT  
 ArgCysGluArgValAspLeuPheTyrLeuArgGlyAspArgGlyGlnIleLeuValIle  
 80 90  
 670 680 690 700 710 720  
 TGTTTGATAGCAGTTATGGTAGTTTTTATTATTTTGGTCATCGGTGTCTGCACATGCTGT  
 CysLeuIleAlaValMetValValPheIleIleLeuValIleGlyValCysThrCysCys  
 100 110

4 / 2 0

F I G . 3 - 2

730 740 750 760 770 780  
CACCTCTTCGGAACGTCGTAAAAGAAAGAAGAAAGAAGAAATGGAACTCTGGGT  
HisProLeuArgLysArgArgLysArgLysLysLysGluGluGluMetGluThrLeuGly  
120 130

790 800 810 820 830 840  
AAAGATATAACTCCTATCAATGAAGATATTGAAGAGACAAATATTGCTTAAAAGGCTATG  
LysAspIleThrProIleAsnGluAspIleGluGluThrAsnIleAlaEnd  
140 147

850 860 870 880 890 900  
AAGTTACCTCCAGGTTGGTGGCAAGCTGCAAAGTGCCTTGCTCATTTGAAATGGACAGA

910 920 930 940 950 960  
ATGTGTCTCAGGAAAAACAGCTAGTAGACATGAATTTTAAATAATGTATTTACTTTTAT

970 980 990 1000 1010 1020  
TTGCAACTTTAGTTTGTGTTATTATTTTTTAATAAGAACATTAATTATATGTATATTGTC

1030 1040 1050 1060 1070 1080  
TAGTAATTGGGAAAAAGCAACTGGTTAGGTAGCAACAACAGAAGGGAAATTTCAATAAC

1090 1100 1110 1120 1130 1140  
CTTTCACTTAAGTATTGTCACCAGGATTACTAGTCAAACAAAAAGAAAGTAGAAAGGA


1150 1160 1170 1180 1190 1200  
GGTTAGGTCTTAGGAATTGAATTAATAATAAAGCTACCATTTATCAAGCATTTACCATGT

1210 1220 1230 1240 1250 1260  
GCTAATAAGTTTGAAATATATTATTTCTTTATTCCTTTCAGCAATCCATGAGATAGCTA

1270  
TTATAATCCTC


5 / 20

FIG. 4




6 / 20

FIG. 5




7 / 20

FIG. 6



8 / 20


FIG.7





9 / 20

FIG. 8



10/ 20

FIG.9A






FIG.9B




11 / 20

FIG.10




12 / 20

FIG.11




13 / 20

FIG.12



14/ 20

FIG.13



1 5 / 2 0

FIG. 14


1 cDNA  
 M D G N S T R S P E T N G L L C G D P E  
 | | | | | | | | | | | | | | |  
 BTC-I M D G N X T R S P E T N G L L X G D P E

21  
 E N C A A T T T Q S K R | | R K G X F S R C P  
 | | | | | | | | | | | | | | |  
 BTC-II R K G X F S R X P

41  
 K Q Y K H Y C I K G R C R F V V A E Q T  
 | | | | | | | | | | | | | | |  
 K Q Y K H Y X I K G R

16/ 20


FIG.15





17 / 20

FIG. 16




18/20

FIG. 17




19/20



20/20

FIG. 19



# INTERNATIONAL SEARCH REPORT

International Application No  
PCT/JP 96/00762

<b>A. CLASSIFICATION OF SUBJECT MATTER</b> IPC 6 C12N15/06 C07K16/22 C12N5/20 G01N33/577 G01N33/68 A61K39/395 C07K1/22 //C07K14/475		
According to International Patent Classification (IPC) or to both national classification and IPC		
<b>B. FIELDS SEARCHED</b> Minimum documentation searched (classification system followed by classification symbols) IPC 6 C12N C07K G01N A61K		
Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched		
Electronic data base consulted during the international search (name of data base and, where practical, search terms used)		
<b>C. DOCUMENTS CONSIDERED TO BE RELEVANT</b>		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	EP,A,0 555 785 (TAKEDA CHEMICAL INDUSTRIES, LTD. & CHILDREN'S MEDICAL CENTER CORP.) 18 August 1993 cited in the application see page 8, line 56 - page 9, line 9 see figure 10 see claims	1-15
A	--- EP,A,0 482 623 (CHILDREN'S MEDICAL CENTER CORP.) 29 April 1992 cited in the application see page 3, line 19 - line 26 see claims --- <div style="text-align: right;">-/--</div>	1-15
<div style="display: flex; justify-content: space-between;"> <span><input checked="" type="checkbox"/> Further documents are listed in the continuation of box C.</span> <span><input checked="" type="checkbox"/> Patent family members are listed in annex.</span> </div>		
<div style="display: flex;"> <div style="flex: 1;"> <p>* Special categories of cited documents :</p> <p>"A" document defining the general state of the art which is not considered to be of particular relevance</p> <p>"E" earlier document but published on or after the international filing date</p> <p>"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</p> <p>"O" document referring to an oral disclosure, use, exhibition or other means</p> <p>"P" document published prior to the international filing date but later than the priority date claimed</p> </div> <div style="flex: 1;"> <p>"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</p> <p>"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone</p> <p>"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.</p> <p>"&amp;" document member of the same patent family</p> </div> </div>		
Date of the actual completion of the international search	Date of mailing of the international search report	
21 June 1996	09.07.96	
Name and mailing address of the ISA European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tel. (+ 31-70) 340-2040, Tx. 31 651 epo nl, Fax: (+ 31-70) 340-3016	Authorized officer  Nooij, F	

## INTERNATIONAL SEARCH REPORT

International Application No

PCT/JP 96/00762

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	<p>BIOCHEMICAL AND BIOPHYSICAL RESEARCH COMMUNICATIONS, vol. 190, no. 3, 15 February 1993, DULUTH, MN, USA, pages 1173-1179, XP002006317 R. SASADA ET AL.: "Cloning and expression of cDNA encoding human betacellulin, a new member of the EGF family." cited in the application see abstract see figure 1</p> <p>---</p>	1-15
A	<p>THE JOURNAL OF BIOLOGICAL CHEMISTRY, vol. 269, no. 13, 1 April 1994, BALTIMORE, MD, USA, pages 9966-9973, XP002006318 T. WATANABE ET AL.: "Recombinant human betacellulin." cited in the application see abstract see table II</p> <p>---</p>	1-15
A	<p>SCIENCE, vol. 259, no. 5101, 12 March 1993, WASHINGTON, DC, USA, pages 1604-1607, XP002006319 Y. SHING ET AL.: "Betacellulin: a mitogen from pancreatic beta cell tumors." cited in the application see figure 2</p> <p>-----</p>	1-15

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/JP 96/00762

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP-A-555785	18-08-93	JP-A- 6087894 CA-A- 2089111	29-03-94 11-08-93
EP-A-482623	29-04-92	US-A- 5229493 AT-T- 122358 CA-A- 2054289 DE-D- 69109615 DE-T- 69109615 JP-A- 4352800 US-A- 5328986	20-07-93 15-05-95 27-04-92 14-06-95 09-11-95 07-12-92 12-07-94